

# Modular Synthesis of Diverse Natural Product-like Macrocycles

Dow, Mark; Marchetti, Fransceco; Abrahams, Katherine; Vaz, Luis; Besra, Gurdyal; Warriner, Stuart

DOI:

[10.1002/chem.201701150](https://doi.org/10.1002/chem.201701150)

License:

None: All rights reserved

Document Version

Peer reviewed version

Citation for published version (Harvard):

Dow, M, Marchetti, F, Abrahams, K, Vaz, L, Besra, G & Warriner, S 2017, 'Modular Synthesis of Diverse Natural Product-like Macrocycles: Discovery of Hits with Antimycobacterial Activity', *Chemistry: A European Journal*.  
<https://doi.org/10.1002/chem.201701150>

[Link to publication on Research at Birmingham portal](#)

## Publisher Rights Statement:

Checked for eligibility: 10/04/2017.

This is the peer reviewed version of the following article: Dow, M., Marchetti, F., Abrahams, K., Vaz, L., Besra, G., Warriner, S. and Nelson, A. (2017), Modular Synthesis of Diverse Natural Product-like Macrocycles: Discovery of Hits with Antimycobacterial Activity. *Chem. Eur. J.*, which has been published in final form at doi:10.1002/chem.201701150. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Self-Archiving.

## General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

## Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.

# CHEMISTRY

## A European Journal

A Journal of



### Accepted Article

**Title:** Modular Synthesis of Diverse Natural Product-like Macrocycles:  
Discovery of Hits with Antimycobacterial Activity

**Authors:** Mark Dow, Fransceco Marchetti, Katherine Abrahams, Luis  
Vaz, Gurdyal Besra, stuart warriner, and Adam Nelson

This manuscript has been accepted after peer review and appears as an Accepted Article online prior to editing, proofing, and formal publication of the final Version of Record (VoR). This work is currently citable by using the Digital Object Identifier (DOI) given below. The VoR will be published online in Early View as soon as possible and may be different to this Accepted Article as a result of editing. Readers should obtain the VoR from the journal website shown below when it is published to ensure accuracy of information. The authors are responsible for the content of this Accepted Article.

**To be cited as:** *Chem. Eur. J.* 10.1002/chem.201701150

**Link to VoR:** <http://dx.doi.org/10.1002/chem.201701150>

Supported by  
**ACES**

WILEY-VCH

# Modular Synthesis of Diverse Natural Product-like Macrocycles: Discovery of Hits with Antimycobacterial Activity


Mark Dow,<sup>[a,b]</sup> Francesco Marchetti,<sup>[a,b]</sup> Katherine A. Abrahams,<sup>[c]</sup> Luis Vaz,<sup>[d]</sup> Gurdyal S. Besra,<sup>[c]</sup> Stuart Warriner<sup>[a,b]</sup> and Adam Nelson<sup>\*[a,b]</sup>

**Abstract:** A modular synthetic approach was developed in which variation of the triplets of building blocks used enabled systematic variation of the macrocyclic scaffolds prepared. The approach was demonstrated in the synthesis of 17 diverse natural product-like macrocyclic scaffolds of varied (12-20-membered) ring size. The biological relevance of the chemical space explored was demonstrated through the discovery of a series of macrocycles with significant antimycobacterial activity.


The remarkable biological functions of macrocyclic natural products have inspired over 100 marketed drugs, predominantly based on macrocyclic peptides and macrolides.<sup>[1,2]</sup> Yet, despite an increasing recognition of their virtues, macrocycles remain an under-represented compound class in drug discovery.<sup>[3]</sup> Macrocycles strike a valuable balance between structural pre-organization, and the flexibility that is needed to target some challenging binding sites.<sup>[1]</sup> In addition, macrocyclisation can increase potency and selectivity dramatically, and improve physicochemical and pharmacokinetic properties.<sup>[1,2,4]</sup> Furthermore, an understanding of molecular properties pertinent to macrocyclic drug discovery is emerging, together with the determinants of cell permeability.<sup>[2,5]</sup> Although macrocycles that target extended binding sites (e.g. in viral proteases and polymerases) typically lie outside conventional drug-like chemical space, some bioactive macrocycles (e.g. some kinase inhibitors) can have similar properties to other small molecule drugs.<sup>[2]</sup>

The discovery of bioactive macrocycles has been hampered by the historic uneven and unsystematic exploration of the relevant chemical space.<sup>[6]</sup> As a result, there is a paucity of macrocycles in established small molecule screening collections. Recently, approaches<sup>[7]</sup> have been developed for the synthesis of diverse non-peptidic macrocycles, for example from building

blocks,<sup>[8]</sup> by ring expansion,<sup>[9]</sup> and using DNA-templated methods.<sup>[10]</sup> In addition, genetically-encoded approaches can enable discovery of bioactive macrocyclic peptides.<sup>[11]</sup> The value of approaches to explore the properties of diverse macrocycles is reflected in the discovery of macrocyclic antimalarials,<sup>[12]</sup> angiogenesis inhibitors,<sup>[13]</sup> histone deacetylase inhibitors<sup>[14]</sup> and blockers of sonic hedgehog signalling.<sup>[15]</sup>



**Scheme 1.** Envisaged approach to skeletally-diverse macrocycles. <sup>F</sup>DIPEO, diisopropyl (3,3,4,4,5,5,6,6,7,7,8,8,9,9,10,10,10-heptafluorodecyl)silyl; Ns, *o*-nitro phenylsulfonyle.



**Figure 1.** Structures of building blocks. Panel A: Initiating building blocks. Panel B: Propagating building blocks. Panel C: Terminating building blocks. Ar, 2,4-dimethoxyphenyl.

Previously, we reported the diversity-oriented synthesis of natural product-like molecules based on >80 diverse molecular scaffolds.<sup>[16]</sup> Here, the product scaffolds depended on both the order and nature of the attachment of pairs of building blocks to a fluorine-tagged linker. We envisaged that the exploitation of triplets of building blocks would enable extension to the synthesis of skeletally-diverse macrocycles (Scheme 1). Thus,

[a] Dr Mark Dow, Dr Francesco Marchetti, Dr Stuart Warriner and Prof. Adam Nelson  
School of Chemistry  
University of Leeds  
Leeds, LS2 9JT, UK  
E-mail: a.s.nelson@leeds.ac.uk

[b] Dr Mark Dow, Dr Francesco Marchetti, Dr Stuart Warriner and Prof. Adam Nelson  
Astbury Centre for Structural Molecular Biology  
University of Leeds  
Leeds, LS2 9JT, UK


[c] Dr Katherine A. Abrahams and Prof. Gurdyal S. Besra  
School of Biosciences  
University of Birmingham  
Edgbaston, Birmingham, B15 2TT, UK

[d] Luis Vaz,  
AstraZeneca,  
Charter Way,  
Macclesfield, SK10 2NA, UK

Supporting information for this article is given via a link at the end of the document.

iterative attachment of propagating and terminating building blocks (Figure 1) to a fluoros-tagged initiating building block **1** ( $\rightarrow$  **2**) would be followed by macrocyclisation ( $\rightarrow$  **3**). We recognised that the fluoros–solid phase extraction<sup>[17]</sup> (F–SPE) could enable removal of excess reactants at each stage of the synthesis, and facilitate decoration of the product macrocycle.

Initially, propagating building blocks were attached to the fluoros-tagged initiating building blocks **1**, **4** and *ent*-**4**, followed by deacetylation (Scheme 2). In each case, the fluoros-tagged products (**13–17**) were isolated after F–SPE.




**Scheme 2.** Attachment of propagating building blocks to fluoros-tagged initiating building blocks. Methods: A: PPh<sub>3</sub>, DEAD, THF then F–SPE; B: NH<sub>3</sub> in MeOH. <sup>a</sup>Mass recovery (and purity determined by 500 MHz <sup>1</sup>H NMR spectroscopy) after purification by F–SPE alone.

Next, we determined the competence of alternative classes of terminating building block. Thus, the *N*-*o*-nitrophenylsulfonyl amide **8**, the *N*-*o*-nitrophenylsulfonyl carbamate **9** and the trifluoromethanesulfonamide **11** were appended to appropriate substrates (**13**, **15** and **17**) (Scheme 3). The ring-closing metatheses<sup>[18]</sup> of the dienes **2** and **18** (5 mM) proceeded smoothly with Hoveyda–Grubbs second generation catalyst in methyl *tert*-butyl ether (MTBE) at 55 °C to give the corresponding macrocycles **3** and **19**. However, the clean ring-closing metathesis of **20** required 10 mol% 1,4-benzoquinone<sup>[18b]</sup> in addition to 5 mol% catalyst to give, after treatment with thiophenol and potassium carbonate, the macrocyclic carbamate **21** (R<sup>1</sup>=<sup>F</sup>DIPES; R<sup>2</sup>=R<sup>3</sup>=H) in 34% yield. In a similar vein, treatment of the diene **22** with 2 mol% catalyst and 4 mol% 1,4-benzoquinone gave the corresponding macrocycle **23** (R<sup>1</sup>=<sup>F</sup>DIPES; R<sup>2</sup>=R<sup>3</sup>=H) in 84% yield.

We then investigated the deprotection of the exemplar macrocycles. The attempted deprotection of the macrocycle **3** did not proceed smoothly. Treatment of **3** with thiophenol and potassium carbonate, followed by F–SPE eluting with MeOH–water and then methanol, resulted in overall methanolysis of the macrolactam; presumably, the Ns group promotes thiolysis of the macrolactam to give a thioester which is subsequently methanolised. This unwanted reaction did not plague the deprotection of the corresponding diene **2**, nor the larger macrolactam **19**, and may be attributed to the strain of the


macrocyclic ring system of **3**. In contrast, desulfonylation of the macrocycles **21** and **23** proceeded smoothly. It was decided that *N*-*o*-nitrophenylsulfonyl amides would not be exploited as terminating building blocks in the synthesis of the macrocycle library.



**Scheme 3.** Attachment of terminating building blocks and subsequent ring-closing metathesis. Methods: A: PPh<sub>3</sub>, DEAD, CH<sub>2</sub>Cl<sub>2</sub> then F–SPE; C: 1 mol% Hoveyda–Grubbs second generation catalyst, MTBE, 55 °C then P(CH<sub>2</sub>OH)<sub>3</sub>, Et<sub>3</sub>N, silica; D: PhSH, K<sub>2</sub>CO<sub>3</sub>, DMF. <sup>a</sup>Mass recovery (and purity determined by 500 MHz <sup>1</sup>H NMR spectroscopy) after purification by F–SPE alone. <sup>b</sup>2 mol% catalyst was used. <sup>c</sup>5 mol% catalyst and 10 mol% 1,4-benzoquinone was used. <sup>d</sup>2 mol% catalyst and 4 mol% 1,4-benzoquinone was used. Ar, 2,4-dimethoxyphenyl.

The synthesis of a wide range of macrocyclic scaffolds is summarised in Table 1 and Figure 2. The cyclisation precursors were prepared by treatment of a fluoros-tagged substrate (**13–17**) and a terminating building block (**9**, **10**, **11** or **12**) with triphenylphosphine and DEAD: the fluoros-tagged product was generally purified by F–SPE and used without further purification. The cyclisation precursors were treated with Hoveyda–Grubbs second generation catalyst (typically 2 mol%), usually in the presence of 1,4-benzoquinone. In general, the cyclisation reactions proceeded smoothly, and macrocyclic

products were isolated after column chromatography. The stereoselectivity of the metathesis reaction was often high, and, in many other cases, the geometric isomers were separable after subsequent desulfonylation.



**Figure 2.** Structures of the natural product-like macrocycles prepared by ring-closing metathesis ( $R^1 = \text{FIPES}$ ;  $R^2 = R^3 = \text{Ns}$ ) (see Table 1 for details). Subsequently, the products were deprotected and decorated to give final compounds ( $R^1 = R^3 = \text{H}$ ;  $R^2 = \text{H}$ , cyclopropylcarbonyl, 1-methyl-imidazol-4-yl sulfonyl or 3-pyridylaminocarbonyl).

Crucially, it was demonstrated that a modular synthesis of macrocyclic scaffolds was possible. By variation of the building blocks used, 17 different macrocyclic scaffolds were prepared with ring sizes varying from 12 to 20. The distributions of macrocycle ring sizes and natural product likeness scores<sup>[19]</sup> of the deprotected scaffolds were compared with those of 2435 commercially-available<sup>[20]</sup> 12- to 20-membered macrocycles (Figure 3). The 17 macrocyclic scaffolds all had positive natural product-likeness scores that reflect their local natural product-like structural features.

The macrocyclic scaffolds were subsequently deprotected and decorated (see Scheme 4 for an


example). For example, the macrocycle **32** ( $R^1 = \text{FIPES}$ ;  $R^2 = R^3 = \text{Ns}$ ) was treated with thiophenol and potassium carbonate in DMF, and the resulting product was purified by F-SPE. The product was then desilylated ( $\rightarrow$  **39c**), or decorated prior to desilylation (e.g.  $\rightarrow$  **39a** or **39b**). In total, 66 products based on the 17 macrocyclic scaffolds were prepared.




**Table 1.** Synthesis of natural product-like macrocycles (see Figure 2; Products had R<sup>1</sup>=<sup>F</sup>DIPES; R<sup>2</sup>, R<sup>3</sup> = Ns; R<sup>4</sup> = TBS unless stated).

Building blocks	Attachment of terminating building block <sup>[a]</sup>	Metathesis <sup>[c]</sup>		
	Mass recovery (purity) <sup>[b]</sup>	Product	Yield <sup>[d]</sup>	E:Z <sup>[e]</sup>
13, 9	66 (95)	24 <sup>[f]</sup>	33 <sup>[g]</sup>	60:40
13, 10	77 (93)	25 <sup>[h]</sup>	56	65:35
13, 11	57 <sup>[d]</sup>	23 <sup>[f]</sup>	84 <sup>[g]</sup>	30:70
13, 12	93 (>95)	26	76	>98:<2
14, 9	>98 (59)	27	56	60:40
14, 10	76 (94)	28 <sup>[h]</sup>	88	[i]
14, 11	93 (95)	29 <sup>[h]</sup>	78	15:85
14, 12	>98 (83)	30 <sup>[h]</sup>	47	[i]
15, 9	90 (93)	21 <sup>[f]</sup>	34 <sup>[g]</sup>	65:35
15, 10	70 <sup>[d]</sup>	31	60	>98:<2
16, 9	97 <sup>[d]</sup>	32	88	>98:<2
16, 10	95 <sup>[d]</sup>	33	75	>98:<2
16, 11	93 <sup>[d]</sup>	34	61	>98:<2
16, 12	>98 <sup>[d]</sup>	35	98	70:30
17, 9	92 <sup>[d]</sup>	36	76	25:75
17, 10	90 <sup>[d]</sup>	37	78	70:30
17, 12	99 <sup>[d]</sup>	38 <sup>[i]</sup>	55 (E); 15 (Z)	

[a] Method A: PPh<sub>3</sub>, DEAD, CH<sub>2</sub>Cl<sub>2</sub> then F-SPE. [b] Mass recovery (and purity determined by 500 MHz <sup>1</sup>H NMR spectroscopy) after purification by F-SPE alone. [c] Method C: 2 or 5 mol% Hoveyda-Grubbs second generation catalyst, 0, 4 or 10 mol% 1,4-benzoquinone, MTBE, 55 °C then P(CH<sub>2</sub>OH)<sub>3</sub>, Et<sub>3</sub>N, silica (see Supporting Information). [d] Yield after column chromatography. [e] Determined by 500 MHz <sup>1</sup>H NMR spectroscopy. [f] The product had R<sup>1</sup>=<sup>F</sup>DIPES; R<sup>2</sup>=R<sup>3</sup>=H after desulfonylation. [g] Yield over two steps after desulfonylation with PhSH, K<sub>2</sub>CO<sub>3</sub>. [h] The geometric isomers were separable after subsequent desulfonylation. [i] Not determined at this stage; after desulfonylation, **28** and **30** were separable geometric isomers. [j] The geometrical isomers were separable.



**Figure 3.** Macrocycle size and natural product-likeness scores of the 17 deprotected macrocyclic scaffolds prepared (black dots) and 2435 commercially-available macrocycles (blue dots).





**Scheme 4.** Exemplar decoration of a macrocyclic scaffold. Method D: PhSH, K<sub>2</sub>CO<sub>3</sub>; Method E: urea or sulfonamide formation; Method F: 50% aq. HF, CH<sub>2</sub>Cl<sub>2</sub>-MeCN then Me<sub>3</sub>SiOMe.

To demonstrate biological relevance, the antimycobacterial activity of the 66 decorated macrocyclic products was assessed using *Mycobacterium bovis* BCG, a model organism for *Mycobacterium tuberculosis* (Supporting Information). In each case, the bacterium was cultured in the presence of 20 μM macrocycle, and cell viability was assessed by the ability of metabolically-active cells to reduce non-fluorescent resazurin to fluorescent resorufin.<sup>[21]</sup> Under these conditions, the macrocycle **39a** had significant activity, causing 80% growth inhibition.<sup>[22]</sup> The IC<sub>50</sub> of the macrocycle **39a** (IC<sub>50</sub> = 11 μM) was determined, together with those of a range of analogues: **39b** and **39c** which bear different substituents (IC<sub>50</sub> = 35 and 30 μM respectively); the isomeric *meta* cyclophane **40a** (IC<sub>50</sub> = 21 μM); and the ring-contracted analogue **41a** (IC<sub>50</sub> = 24 μM) (Figure 4). Taken together, the initial single concentration and subsequent concentration-dependent activity determinations suggest that both the macrocyclic scaffold and the substituents are important for biological function. Hits from such phenotypic screens are highly valuable since they can facilitate the identification of new targets for antimycobacterial drug discovery.<sup>[23]</sup> Crucially, the macrocycle **39a** is highly distinctive from the TB box set secured

from a high-throughput screen of GSK's collection:<sup>[24]</sup> none of the 177 compounds in the box set contains a macrocyclic ring.

A



**Figure 4.** Effect of macrocycles on the viability of *M. bovis* BCG. Panel A: Dose-dependent activity of the macrocycle **39a** and four analogues. Panel B: Antimycobacterial activity of selected macrocycles.

In summary, we have developed a modular approach that was exploited in the synthesis of 17 diverse natural product-like macrocyclic scaffolds. Through variation of the triplets of building blocks used, systematic variation of the scaffolds and macrocyclic ring size (12-20-membered) was possible. The

biological relevance of chemical space explored was demonstrated through discovery of a series of distinctive macrocycles with significant antimycobacterial activity.

## Acknowledgements

We thank EPSRC (EP/F43503/1; EP/N025652/1) and AstraZeneca for funding. GSB acknowledges support in the form of a Personal Research Chair from Mr James Bardrick, a Royal Society Wolfson Research Merit Award, the Medical Research Council (MR/K012118/1) and the Wellcome Trust (081569/Z/06/Z).

**Keywords:** diversity-oriented synthesis • macrocycles • antimycobacterials


- [1] E. M. Driggers, S. P. Hale, N. K. Terrett, *Nat. Rev. Drug. Discov.* **2008**, 7, 608-624.
- [2] F. Giordanetto, J. Kihlberg, *J. Med. Chem.* **2014**, 57, 278-295.
- [3] (a) R. D. Taylor, M. MacCoss, A. D. G. Lawson, *J. Med. Chem.* **2014**, 57, 5845-5859; (b) H. Waldmann, E. Valeur, S. M. Guéret, H. Adihou, R. Gopalakrishnan, M. Lemurell, T. N. Grossmann, A. T. Plowright, *Angew. Chem. Int. Ed.* **2017**, DOI: 10.1022/anie.201611914.
- [4] For examples, see: (a) C. J. Dinsmore *et al.*, *J. Am. Chem. Soc.* **2001**, 123, 2107-2108; (b) Z.-F. Tao *et al.*, *J. Med. Chem.* **2007**, 50, 1514-1527; (c) R. J. Cherney *et al.*, *J. Med. Chem.* **1998**, 41, 1749-1751; (d) J. R. Corte *et al.*, *J. Med. Chem.* **2017**, DOI:10.1021/acs.jmedchem.6b01460.
- [5] (a) B. Over *et al.*, *Nat. Chem. Biol.* **2016**, 12, 1065-1074; (b) C. R. Pye *et al.*, *J. Med. Chem.* **2017**, DOI:10.1021/acs.jmedchem.6b01483.
- [6] A. H. Lipkus, Q. Yuan, K. A. Lucas, S. A. Funk, W. F. Bartlett III, R. J. Schenk, A. J. Trippie, *J. Org. Chem.* **2008**, 73, 4443-4451.
- [7] S. Collins, S. Bartlett, F. Nie, H. F. Sore, D. R. Spring, *Synthesis* **2016**, 48, 1457-1473.
- [8] For examples, see: (a) E. Comer, H. Liu, A. Joliton, A. Clabaut, C. Johnson, L. B. Akella, L. A. Marcaurelle, *Proc. Natl. Acad. Sci. USA* **2011**, 108, 6751-6756; (b) A. Grossmann, S. Bartlett, M. Janacek, J. T. Hodgkinson, D. R. Spring, *Angew. Chem. Int. Ed.* **2014**, 53, 13093-13097; (c) H. S. G. Beckmann, F. Nie, C. E. Hagerman, H. Johansson, Y. S. Tan, D. Wilcke, D. R. Spring, *Nat. Chem.* **2013**, 5, 861-867; (d) F. Nie, D. L. Kunciw, D. Wilcke, J. E. Stokes, W. R. J. D. Galloway, S. Bartlett, H. F. Sore, D. R. Spring, *Angew. Chem. Int. Ed.* **2016**, 55, 11139-11143.
- [9] For examples, see: (a) F. Kopp, C. F. Stratton, L. B. Akella, D. S. Tan, *Nat. Chem. Biol.* **2012**, 8, 358-365; (b) C. Kitsiou, J. J. Hindes, P. l'Anson, P. Jackson, T. C. Wilson, E. K. Daly, H. R. Felstead, P. Hearnshaw, W. P. Unsworth, *Angew. Chem. Int. Ed.* **2015**, 54, 15794-15798.
- [10] For a review, see: W. H. Connors, S. P. Hale, N. K. Terrett, *Curr. Opin. Chem. Biol.* **2015**, 26, 42-47.
- [11] For a review, see: K. R. Lennard, A. Tavassoli, *Chem. Eur. J.* **2014**, 20, 10608-10614.
- [12] E. Comer *et al.*, *J. Med. Chem.* **2014**, 57, 8496-8502.
- [13] M. Aeluri, C. Pramanik, L. Chetia, N. K. Mallurwar, S. Balasubramanian, G. Chandrasekar, S. S. Kitambi, P. Arya, *Org. Lett.* **2013**, 15, 436-439.
- [14] L. A. Marcaurelle *et al.*, *J. Am. Chem. Soc.* **2010**, 132, 16962-16976.
- [15] B. Z. Stanton, L. F. Peng, N. Maloof, K. Nakai, X. Wang, J. L. Duffner, K. M. Taveras, J. M. Hyman, S. W. Lee, A. N. Koehler, J. K. Chen, J. L. Fox, A. Mandinova, S. L. Schreiber, *Nat. Chem. Biol.* **2009**, 5, 154-156.
- [16] D. Morton, S. Leach, C. Cordier, S. Warriner, A. Nelson, *Angew. Chem. Int. Ed.* **2009**, 48, 104-109.

- [17] *Handbook of Fluorous Chemistry* (Eds.: J. A. Gladysz, D. P. Curran, I. T. Horváth), Wiley-VCH, Weinheim, **2005**.
- [18] (a) K. M. Kuhn, T. M. Champagne, S. H. Hong, W.-H. Wei, A. Nickel, C. W. Lee, S. C. Virgil, R. H. Grubbs, R. L. Pederson, *Org. Lett.* **2010**, *12*, 984-987; (b) S. H. Hong, D. P. Sanders, C. W. Lee, R. H. Grubbs, *J. Am. Chem. Soc.* **2005**, *127*, 17160-17161.
- [19] P. Ertl, S. Roggo, A. Schuffenhauer, *J. Chem. Inf. Model.*, **2008**, *48*, 68-74. Natural product likeness scores were calculated using the implementation in RDKit v2015.09.2 (Greg Landrum; Open Source Cheminformatics; <http://www.rdkit.org>; last accessed 27-Apr-2016).
- [20] J. J. Irwin, T. Sterling, M. M. Mysinger, E. S. Bolstad, R. G. Coleman, *J. Chem. Inf. Model.* **2012**, *52*, 1757-1768.
- [21] (a) S. G. Franzblau, R. S. Witzig, J. C. McLaughlin, P. Torres, G. Madico, A. Hernandez, M. T. Degnan, M. B. Cook, V. K. Quenzer, R. M. Ferguson, R. H. Gilman, *J. Clin. Microbiol.* **1998**, *36*, 362-366; (b) S. M. Batt *et al*, *ACS Infect. Dis.* **2015**, *1*, 615-626.
- [22] In this preliminary single concentration screen, the macrocycle **39a** was the only compound whose activity was more than two standard deviations (>22% growth inhibition) from the mean.
- [23] For examples, see: (a) K. A. Abrahams *et al*, *Nat. Commun.* **2016**, *7*, 12581; (b) J. A. G. Cox *et al*, *Nature Microbiol.* **2016**, *1*, 15006.
- [24] L. Ballell *et al*, *ChemMedChem* **2013**, *8*, 313-321.



## Entry for the Table of Contents

## COMMUNICATION



A modular approach enabled the synthesis of 17 diverse natural product-like macrocyclic scaffolds, and the discovery of a distinctive macrocycles with antimycobacterial activity.

Mark Dow, Francesco Marchetti,  
Katherine A. Abrahams, Luis Vaz,  
Gurdial S. Besra, Stuart Warriner and  
Adam Nelson\*

Page No. – Page No.

**Modular Synthesis of Diverse Natural  
Product-like Macrocycles: Discovery  
of Hits with Antimycobacterial  
Activity**